

HUMAN FGF GENE AND GENE EXPRESSION PRODUCTS

REFERENCE TO RELATED APPLICATIONS

This application is a continuation of provisional U.S. Patent Application Serial No. 60/077,411, filed March 9, 1998, and U.S. Patent Application Serial No. 60/083,553, filed April 29, 1998.

INSAI

TECHNICAL FIELD

The present invention relates to novel nucleic acid sequences encoding a member of the fibroblast growth factor (FGF) family, and to polypeptides encoded by the nucleic acid sequences.

BACKGROUND OF THE INVENTION

This invention provides a gene of the fibroblast growth factor (FGF) family and protein encoded by the gene. The invention is directed to mRNA expressed by human cells, polynucleotides having coding regions corresponding to the mRNA, protein products of the polynucleotides and polypeptide products of mRNA, and biological function of the polypeptides and proteins.

At least 17 members of the FGF family have been identified, and are discussed in, for example, Ohuchi et al., *Development* 124:2235-2244 (1994); Ghosh et al., *Cell Growth and Differentiation* 7:1425-1434 (1996); Gemel et al., *Genomics* 35:253-257 (1996); Crossley et al., *Development* 121:439-451 (1995); and Crossley et al., *Cell* 84:127-136 (1996).

SUMMARY OF THE INVENTION

The present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising at least eight contiguous nucleotides from nucleotide 550 to 1170 of SEQ ID NO:4;
- (b) a polynucleotide that encodes a variant of the polypeptide encoded by (a); and

- (c) a polynucleotide encoding a protein expressed by a polynucleotide having the sequence of SEQ ID NO:4.

The invention further provides for the use of the isolated polynucleotides or fragments thereof as diagnostic probes or as primers.

The present invention also provides a composition comprising a polypeptide, wherein said polypeptide is selected from the group consisting of:

- (a) a polypeptide comprising at least 6 contiguous amino acids encoded by SEQ ID NO:4 in the region of nucleotide 550 to 1170;
- (b) a polypeptide encoded by a polynucleotide comprising SEQ ID NO:4; and
- (c) a variant of the protein (a) or (b).

In certain preferred embodiments of the invention, the polynucleotide is operably linked to an expression control sequence. The invention further provides a host cell, including bacterial, yeast, insect and mammalian cells, transformed with the polynucleotide sequence. The invention also provides full-length cDNA and full-length polynucleotides corresponding to SEQ ID NO:4.

Protein and polypeptide compositions of the invention may further comprise a pharmaceutically acceptable carrier. Compositions comprising an antibody that specifically reacts with such protein or polypeptide are also provided by the present invention.

The invention provides methods of using the FGF of the invention for the isolation, regeneration, proliferation, and differentiation of mammalian multipotent neural stem cells, progenitor cells and progeny.

In a preferred embodiment, primary central (CNS) and peripheral nervous system (PNS) cells when treated with FGF 98 of the invention proliferate, have at least a limited self regeneration capacity, and can undergo lineage restriction in response to the local environment.

In a further embodiment, cells produced by treatment with FGF 98 are used to screen drugs such as small molecules, and growth factors, which may affect development, differentiation, survival and or function of CNS and PNS derived neurons and glia.

The invention also provides for the production of large amounts of otherwise minor cell populations of cells to be used for generation of cDNA libraries for the isolation of rare molecules expressed in the precursors cells or progeny; cells produced by treatment may directly express growth factors or other molecules, and conditioned media is screened in assays for novel activities.

The invention further provides for the isolation, self-renewal and survival of mammalian neural stem cells and the differentiation of their progeny.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1. Western analysis of FGF 98 produced by SF-9 cells infected with different clones of baculovirus expressing FGF 98 (Clone 1 to 6). The highest producers were Clones 2 and 4.

Figure 2. (A) Elution profile of the CM of SF-9 cells expressing FGF 98 on a Heparin Sepharose (HS) affinity column. Figure 2(B) shows the OD profile of the 0.5M, 1M and 2M NaCl (● - ●) fractions together with the bioactivity profile of the various fractions (○) when tested on granulosa cells.

Figure 3. (A) shows SDS-PAGE analysis of the 2M NaCl fractions followed by Coomassie blue staining, indicating the presence of a single band with a MW of 28 kDa in the peak fractions of the 2M NaCl eluate (frac. 13, 14). Figure 3(B) shows Western analysis using an anti-peptide FGF 98 antibody of the HS fractions shown in Figure 2A. A single band with a MW of 28 kD is seen in fractions 12 to 16 corresponding to the 2M NaCl fractions.

Figure 4. The active HS fractions (Fractions 2-16) were loaded onto a C4 Vydac column (25x0.46 cm, 5 µm particle size, 300 Å pore size) equilibrated in 0.1% (v/v) TFA, 20% acetonitrile. Protein was eluted with a 115 min. gradient of 20-100% acetonitrile in 0.1% TFA at a flow rate of 0.6 ml/min. at room temperature. Fractions of 1.2 ml were collected. Aliquots of each fraction were diluted 1 to 10 with 0.2% gelatin in PBS and bioassayed on granulosa cells. A major peak of protein closely associated with a small peak could be observed. Both protein peaks were bioactive.

Figure 5. (A) SDS-PAGE of the RP-HPLC fractions shown in Figure 4. When stained by Coomassie blue, both Fractions 22 and 23 gave a single band with a MW of

28 kDa. Figure 5(B). When analyzed by Western using an FGF 98 anti-peptide antibody both Fractions 22 and 23 were recognized as a single band with a MW of 28 kDa.

Figure 6. Heparin Sepharose chromatography of a 3 liter batch of medium conditioned by SF-9 cells expressing FGF 98. The CM was adjusted to 0.5M NaCl and pH to 7.3. It was applied on a column of Heparin Sepharose equilibrated in 0.5M NaCl 10 mM Tris pH 7.2 on sequentially eluted with 1M and 2M NaCl (A, B).

Figure 7. (A) SDS-PAGE of the 1M and 2M NaCl fractions of Figure 6 followed by Coomassie blue staining. Figure 7(B). Western analysis of the 1M and 2M NaCl fractions of Figure 6 using a specific FGF 98 antipeptide antibody.

Figure 8. Mono S ion-exchange chromatography of the HS-purified and active fractions. The bioactive fractions eluted from HS were affinity pooled and diluted 10-fold with 10 mM Tris pH 7.2. Using a 50 ml Super loop, the sample was then applied on a Mono S HR 5/5 column equilibrated in 10 mM Tris pH 7.2-0.2 M NaCl at room temperature. The column was eluted with a multilinear gradient of NaCl (0.2M to 2M NaCl). Absorbency was monitored at 280 nm. Flow rate was 1 ml/min and 1 ml fractions were collected. Aliquots of each fraction were diluted 1 to 100 in 0.2% gelatin in PBS, and 10 μ l aliquots were bioassayed on granulosa cells in 12 well cluster plates as described.

Figure 9. Comparison of the biological activity for granulosa cells of FGF 98 (●) and basic FGF(0).

Figure 10 is a graph showing the effect of FGF 98 on neural stem cell proliferation. Dotted bars represent 6 days *in vitro* (DIV) and solid bars represent 21 days *in vitro* (DIV). The treatment consisted of no additions (1), bFGF (2), or FGF 98 of the invention (3).

Figure 11. Panel A is a photograph of bFGF-treated rat primary hippocampal cell cultures, showing some nestin expression. Panel B shows a separate culture of cells treated with FGF 98 of the invention. More extensive nestin expression is seen.

Figure 12. Panel A shows FGF 98 mRNA expression in adult brain stem and cerebellum. Panel B shows a high magnification of FGF 98 mRNA expression in motor neurons of the facial nucleus.

Fig 2A-D.
Fig 3A-B.
Fig 5A-B
Fig 7A-B
Fig 11
Purified
not
labeled

DETAILED DESCRIPTION OF THE INVENTION

Because of their potent activities for promoting growth, proliferation, survival and differentiation of a wide variety of cells and tissue types, FGFs continue to be pursued as therapeutic agents for a number of different indications, including wound healing, such as musculo-skeletal conditions, for example, bone fractures, ligament and tissue repair, tendonitis, bursitis, etc.; skin conditions, for example, burns, cuts, lacerations, bed sores, slow healing ulcers, etc.; tissue protection and repair during myocardial infarction and ischemia, in the treatment of neurological conditions, for example, neuro-degenerative disease and stroke, in the treatment of eye disease, including macular degeneration, and the like.

The fibroblast growth factor (FGF) proteins identified to date belong to a family of signaling molecules that regulate growth and differentiation of a variety of cell types. The significance of FGF proteins to human physiology and pathology relates in part to their key roles in embryogenesis, in blood vessel development and growth, and in bone growth. *In vitro* experiments have demonstrated a role for FGF in regulating cell growth and division of endothelial cells, vascular smooth muscle cells, fibroblasts, and cardiac and skeletal myocytes. Other members of the FGF family and their biological roles are described in Crossley et al., *Development* 121:439-451 (1995); Ohuchi et al., *Development* 124:2235-2244 (1997); Gemel et al., *Genomics* 35:253-257 (1996); and Ghosh et al., *Cell Growth and Differentiation* 7:1425-1434 (1996).

FGF proteins are also significant to human health and disease because of a role in cancer cell growth. For example, FGF-8 was identified as an androgen-induced growth factor in breast and prostate cancer cells. (Tanaka et al., *FEBS Lett.* 363:226-230 (1995) and *P.N.A.S.* 89:8928-8932 (1992)).

A member of the FGF family is described here, wherein the FGF protein is expressed at high levels in adult mouse brain, and plays a protective role in neuronal tissue. A polynucleotide encoding the FGF of the invention has the sequence as shown in SEQ ID NO:1. The polynucleotide was identified as encoding a member of the FGF family by the conserved regions throughout the amino acid sequence and by the regions of homology shared by the polynucleotide and genes encoding known FGF proteins.

The FGF of the invention shares greatest homology with FGF-8, which plays a variety of roles, including outgrowth and patterning in mouse embryo development (Crossley et al., *Development* 121:439-451 (1995), and midbrain development (Crossley et al., *Nature* 380:66-68 (1996). Eight FGF-8 isoforms have been identified in mice, generated by alternative splicing, and may be functionally redundant. FGF-8b is 100% identical in mice and humans. Blunt et al., *J. Biol. Chem.* 272:3733-3738 (1997). FGF-8 and FGF-10 function in a coordinate manner during chick limb development. Ohuchi et al., *Development* 124:2235-2244 (1997).

As the above brief discussion illustrates, the knowledge about the roles played by various FGF proteins continues to grow, but is by far incomplete.

The present invention adds to this knowledge by disclosing that the FGF of SEQ ID NO:5 plays a role in adult mouse brain and has a protective role in neuronal tissue. The FGF 98 increased the number of neuronal cells at six and twenty days *in vitro*. Furthermore, unlike bFGF-treated cells, few FGF 98-treated cells die under the conditions used herein. FGF 98 can be used for *ex vivo* expansion of precursor cells in the nervous system. Transplant or engraftment of the cells can be used as a therapy for diseases in which specific populations of neurons are lost due to degeneration such as, for example, Parkinson's disease and Alzheimer's disease.

The invention therefore is based upon the identification, isolation and sequencing of a new fibroblast growth factor (FGF 98). This growth factor has been found to promote cell survival and, in particular, the survival of neurons.

Reference to FGF 98 herein is intended to be construed to include growth factors of any origin which are substantially homologous to and which are biologically equivalent to the FGF 98 characterized and described herein. Such substantially homologous growth factors may be native to any tissue or species and, similarly, biological activity can be characterized in any of a number of biological assay systems.

The terms "biologically equivalent" are intended to mean that the compositions of the present invention are capable of demonstrating some or all of the same growth properties in a similar fashion, not necessarily to the same degree as the FGF 98 isolated as described herein or recombinantly produced human FGF 98 of the invention.

By "substantially homologous" it is meant that the degree of homology of human FGF 98 to FGF 98 from any species is greater than that between FGF 98 and any previously reported member of the FGF family.

Sequence identity or percent identity is intended to mean the percentage of same residues between two sequences, referenced to human FGF when determining percent identity with non-human FGF 98, referenced to FGF 98 when determining percent identity with non-FGF 98 growth factors, when the two sequences are aligned using the Clustal method (Higgins et al, *Cabios* 8:189-191, 1992) of multiple sequence alignment in the Lasergene biocomputing software (DNASTAR, INC, Madison, WI). In this method, multiple alignments are carried out in a progressive manner, in which larger and larger alignment groups are assembled using similarity scores calculated from a series of pairwise alignments. Optimal sequence alignments are obtained by finding the maximum alignment score, which is the average of all scores between the separate residues in the alignment, determined from a residue weight table representing the probability of a given amino acid change occurring in two related proteins over a given evolutionary interval. Penalties for opening and lengthening gaps in the alignment contribute to the score. The default parameters used with this program are as follows: gap penalty for multiple alignment=10; gap length penalty for multiple alignment=10; k-tuple value in pairwise alignment=1; gap penalty in pairwise alignment=3; window value in pairwise alignment=5; diagonals saved in pairwise alignment=5. The residue weight table used for the alignment program is PAM250 (Dayhoff et al., in *Atlas of Protein Sequence and Structure*, Dayhoff, Ed., NDRF, Washington, Vol. 5, suppl. 3, p. 345, 1978).

Percent conservation is calculated from the above alignment by adding the percentage of identical residues to the percentage of positions at which the two residues represent a conservative substitution (defined as having a log odds value of greater than or equal to 0.3 in the PAM250 residue weight table). Conservation is referenced to human FGF 98 when determining percent conservation with non-human FGF 98, referenced to FGF 98 when determining percent conservation with non-FGF 98 growth factors. Conservative amino acid changes satisfying this requirement are: R-K; E-D, Y-F, L-M; V-I, Q-H.

FGF 98 can also include hybrid and modified forms of FGF 98 including fusion proteins and FGF 98 fragments and hybrid and modified forms in which certain amino acids

have been deleted or replaced and modifications such as where one or more amino acids have been changed to a modified amino acid or unusual amino acid and modifications such as glycosylations so long as the hybrid or modified form retains the biological activity of FGF 98. By retaining the biological activity, it is meant that neuronal survival is promoted, although not necessarily at the same level of potency as that of the FGF 98 isolated as described herein or that of the recombinantly produced human FGF 98.

Also included within the meaning of substantially homologous is any FGF 98 which may be isolated by virtue of cross-reactivity with antibodies to the FGF 98 described herein or whose encoding nucleotide sequences including genomic DNA, mRNA or cDNA may be isolated through hybridization with the complementary sequence of genomic or subgenomic nucleotide sequences or cDNA of the FGF 98 herein or fragments thereof. It will also be appreciated by one skilled in the art that degenerate DNA sequences can encode human FGF 98 and these are also intended to be included within the present invention as are allelic variants of FGF 98.

Growth factors are thought to act at specific receptors. According to the invention, FGF 98 and as yet unknown members of this family of growth factors act through specific receptors having distinct distributions as has been shown for other growth factor families.

A preferred FGF 98 of the present invention has been identified and isolated in purified form as described. Also preferred is FGF 98 prepared by recombinant DNA technology. By "pure form" or "purified form" or "substantially purified form" it is meant that an FGF 98 composition is substantially free of other proteins which are not FGF 98.

Recombinant human FGF 98 may be made by expressing the DNA sequences encoding FGF 98 in a suitable transformed host cell. Using methods well known in the art, the DNA encoding FGF 98 may be linked to an expression vector, transformed into a host cell and conditions established that are suitable for expression of FGF 98 by the transformed cell.

Any suitable expression vector may be employed to produce recombinant human FGF 98 such as expression vectors for use in insect cells. Baculovirus expression systems can also be employed. A preferable method is expression in insect cells, such as Tr5 or Sf9 cells, using baculovirus vector, as described in Example 1.

The present invention includes nucleic acid sequences including sequences that encode human FGF 98. Also included within the scope of this invention are sequences that are substantially the same as the nucleic acid sequences encoding FGF 98. Such substantially the same sequences may, for example, be substituted with codons more readily expressed in a given host cell such as *E. coli* according to well known and standard procedures. Such modified nucleic acid sequences are included within the scope of this invention.

Specific nucleic acid sequences can be modified by those skilled in the art and, thus, all nucleic acid sequences that code for the amino acid sequences of FGF 98 can likewise be so modified. The present invention thus also includes nucleic acid sequence which will hybridize with all such nucleic acid sequences--or complements of the nucleic acid sequences where appropriate--and encode a polypeptide having the cell survival promoting activities disclosed herein. The present invention also includes nucleic acid sequences that encode for polypeptides that have neuronal survival promoting activity and that are recognized by antibodies that bind to FGF 98. A preferred method for raising antibody is described in Example 9.

The present invention also encompasses vectors comprising expression regulatory elements operably linked to any of the nucleic acid sequences included within the scope of the invention. This invention also includes host cells--of any variety--that have been transformed with vectors comprising expression regulatory elements operably linked to any of the nucleic acid sequences included within the scope of the present invention.

Methods are also provided herein for producing FGF 98. Preparation can be by isolation from conditioned medium from a variety of cell types so long as the cell type produces FGF 98. A second and preferred method involves utilization of recombinant methods by isolating or obtaining a nucleic acid sequence encoding FGF 98, cloning the sequence along with appropriate regulatory sequences into suitable vectors and cell types, and expressing the sequence to produce FGF 98.

Although FGF 98 has been described on the basis of its ability to promote the survival of particular cells, this factor will act on other neuronal cell types as well. For example, FGF 98 is shown herein to enhance nestin expression. Thus, it is likely that FGF 98 can act on a wide variety of central and peripheral neuronal cell types.

It is also likely that FGF 98 will act on non-neuronal cells to promote their survival, growth or function. This expectation is based upon the activity of known growth factors. Members of the FGF family act on many cell types of different function and embryologic origin.

The inventors herein have identified several non-neuronal tissues in which FGF 98 is expressed including lung, heart, developing bone, and chondrocytes. This suggests a role for FGF 98 in, for example, maintenance of musculoskeletal function and treatment of related disease; correction of congenital defects; and treatment cardiomyopathies.

The present invention also includes therapeutic or pharmaceutical compositions comprising FGF 98 in an effective amount for treating patients with cellular degeneration and a method comprising administering a therapeutically effective amount of FGF 98. These compositions and methods are useful for treating a number of degenerative diseases. Where the cellular degeneration involves neuronal degeneration, the diseases include, but are not limited to peripheral neuropathy, amyotrophic lateral sclerosis, Alzheimer's disease, Parkinson's disease, Huntington's disease, ischemic stroke, acute brain injury, acute spinal chord injury, nervous system tumors, multiple sclerosis, peripheral nerve trauma or injury, exposure to neurotoxins, metabolic diseases such as diabetes or renal dysfunctions and damage caused by infectious agents. Where the cellular degeneration involves bone marrow cell degeneration, the diseases include, but are not limited to disorders of insufficient blood cells such as, for example, leukopenias including eosinopenia and/or basopenia, lymphopenia, monocytopenia, neutropenia, anemias, thrombocytopenia as well as an insufficiency of stem cells for any of the above. The above cells and tissues can also be treated for depressed function.

The compositions and methods herein can also be useful to prevent degeneration and/or promote survival in other non-neuronal tissues as well. One skilled in the art can readily determine using a variety of assays known in the art for identifying whether FGF 98 would be useful in promoting survival or functioning in a particular cell type, such as cardiac cells.

In certain circumstances, it may be desirable to modulate or decrease the amount of FGF 98 expressed. Thus, in another aspect of the present invention, FGF 98 anti-

sense oligonucleotides can be made and a method utilized for diminishing the level of expression of FGF 98 by a cell comprising administering one or more FGF 98 anti-sense oligonucleotides. By FGF 98 anti-sense oligonucleotides reference is made to oligonucleotides that have a nucleotide sequence that interacts through base pairing with a specific complementary nucleic acid sequence involved in the expression of FGF 98 such that the expression of FGF 98 is reduced. Preferably, the specific nucleic acid sequence involved in the expression of FGF 98 is a genomic DNA molecule or mRNA molecule that encodes FGF 98. This genomic DNA molecule can comprise regulatory regions of the FGF 98 gene, or the coding sequence for mature FGF 98 protein. The term complementary to a nucleotide sequence in the context of FGF 98 antisense oligonucleotides and methods therefor means sufficiently complementary to such a sequence as to allow hybridization to that sequence in a cell, *i.e.*, under physiological conditions. The FGF 98 antisense oligonucleotides preferably comprise a sequence containing from about 8 to about 100 nucleotides and more preferably the FGF 98 antisense oligonucleotides comprise from about 15 to about 30 nucleotides. The FGF 98 antisense oligonucleotides can also contain a variety of modifications that confer resistance to nucleolytic degradation such as, for example, modified internucleoside linkages (Uhlmann and Peyman, *Chemical Reviews* 90:543-548 1990; Schneider and Banner, *Tetrahedron Lett.* 31:335, 1990 which are incorporated by reference), modified nucleic acid bases and/or sugars and the like.

The therapeutic or pharmaceutical compositions of the present invention can be administered by any suitable route known in the art including for example intravenous, subcutaneous, intramuscular, transdermal, intrathecal or intracerebral. Administration can be either rapid as by injection or over a period of time as by slow infusion or administration of slow release formulation. For treating tissues in the central nervous system, administration can be by injection or infusion into the cerebrospinal fluid (CSF). When it is intended that FGF 98 be administered to cells in the central nervous system, administration can be with one or more agents capable of promoting penetration of FGF 98 across the blood-brain barrier.

FGF 98 can also be linked or conjugated with agents that provide desirable pharmaceutical or pharmacodynamic properties. For example, FGF 98 can be coupled to any substance known in the art to promote penetration or transport across the blood-brain barrier

such as an antibody to the transferrin receptor, and administered by intravenous injection (See, for example, Friden et al., *Science* 259:373-377, 1993 which is incorporated by reference). Furthermore, FGF 98 can be stably linked to a polymer such as polyethylene glycol to obtain desirable properties of solubility, stability, half-life and other pharmaceutically advantageous properties. (See, for example, Davis et al., *Enzyme Eng.* 4:169-73, 1978; Burnham, *Am. J. Hosp. Pharm.* 51:210-218, 1994 which are incorporated by reference).

The compositions are usually employed in the form of pharmaceutical preparations. Such preparations are made in a manner well known in the pharmaceutical art. One preferred preparation utilizes a vehicle of physiological saline solution, but it is contemplated that other pharmaceutically acceptable carriers such as physiological concentrations of other non-toxic salts, five percent aqueous glucose solution, sterile water or the like may also be used. It may also be desirable that a suitable buffer be present in the composition. Such solutions can, if desired, be lyophilized and stored in a sterile ampoule ready for reconstitution by the addition of sterile water for ready injection. The primary solvent can be aqueous or alternatively non-aqueous. FGF 98 can also be incorporated into a solid or semi-solid biologically compatible matrix which can be implanted into tissues requiring treatment.

The carrier can also contain other pharmaceutically-acceptable excipients for modifying or maintaining the pH, osmolarity, viscosity, clarity, color, sterility, stability, rate of dissolution, or odor of the formulation. Similarly, the carrier may contain still other pharmaceutically-acceptable excipients for modifying or maintaining release or absorption or penetration across the blood-brain barrier. Such excipients are those substances usually and customarily employed to formulate dosages for parenteral administration in either unit dosage or multi-dose form or for direct infusion into the cerebrospinal fluid by continuous or periodic infusion.

Dose administration can be repeated depending upon the pharmacokinetic parameters of the dosage formulation and the route of administration used.

It is also contemplated that certain formulations containing FGF 98 are to be administered orally. Such formulations are preferably encapsulated and formulated with suitable carriers in solid dosage forms. Some examples of suitable carriers, excipients, and

diluents include lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, gelatin, syrup, methyl cellulose, methyl- and propylhydroxybenzoates, talc, magnesium, stearate, water, mineral oil, and the like. The formulations can additionally include lubricating agents, wetting agents, emulsifying and suspending agents, preserving agents, sweetening agents or flavoring agents. The compositions may be formulated so as to provide rapid, sustained, or delayed release of the active ingredients after administration to the patient by employing procedures well known in the art. The formulations can also contain substances that diminish proteolytic degradation and promote absorption such as, for example, surface active agents.

The specific dose is calculated according to the approximate body weight or body surface area of the patient or the volume of body space to be occupied. The dose will also be calculated dependent upon the particular route of administration selected. Further refinement of the calculations necessary to determine the appropriate dosage for treatment is routinely made by those of ordinary skill in the art. Such calculations can be made without undue experimentation by one skilled in the art in light of the activity disclosed herein in assay preparations of target cells. Exact dosages are determined in conjunction with standard dose-response studies. It will be understood that the amount of the composition actually administered will be determined by a practitioner, in the light of the relevant circumstances including the condition or conditions to be treated, the choice of composition to be administered, the age, weight, and response of the individual patient, the severity of the patient's symptoms, and the chosen route of administration.

In one embodiment of this invention, FGF 98 may be therapeutically administered by implanting into patients vectors or cells capable of producing a biologically-active form of FGF 98 or a precursor of FGF 98, *i.e.*, a molecule that can be readily converted to a biological-active form of FGF 98 by the body. In one approach cells that secrete FGF 98 may be encapsulated into semipermeable membranes for implantation into a patient. The cells can be cells that normally express FGF 98 or a precursor thereof or the cells can be transformed to express FGF 98 or a precursor thereof. It is preferred that the cell be of human origin and that the FGF 98 be human FGF 98 when the patient is human. However, the

formulations and methods herein can be used for veterinary as well as human applications and the term "patient" as used herein is intended to include human and veterinary patients.

Cells can be grown *ex vivo* for use in transplantation or engraftment into patients (Muench et al., *Leuk. & Lymph.* 16:1-11, 1994 which is incorporated by reference). In another embodiment of the present invention, FGF 98 is used to promote the *ex vivo* expansion of a cells for transplantation or engraftment. Current methods have used bioreactor culture systems containing factors such as erythropoietin, colony stimulating factors, stem cell factor, and interleukins to expand hematopoietic progenitor cells for erythrocytes, monocytes, neutrophils, and lymphocytes (Verfaillie, *Stem Cells* 12:466-476, 1994 which is incorporated by reference). These stem cells can be isolated from the marrow of human donors, from human peripheral blood, or from umbilical cord blood cells. The expanded blood cells are used to treat patients who lack these cells as a result of specific disease conditions or as a result of high dose chemotherapy for treatment of malignancy (George, *Stem Cells* 12(Suppl 1):249-255, 1994 which is incorporated by reference). In the case of cell transplant after chemotherapy, autologous transplants can be performed by removing bone marrow cells before chemotherapy, expanding the cells *ex vivo* using methods that also function to purge malignant cells, and transplanting the expanded cells back into the patient following chemotherapy (for review, see Rummel and Van Zant, *J. Hematotherapy* 3:213-218, 1994 which is incorporated by reference). Since FGF 98 is expressed in the developing animal in blood, bone marrow and liver, tissues where proliferation and differentiation of progenitor cells occur, it is believed that FGF 98 can function to regulate the proliferation of hematopoietic stem cells and the differentiation of mature hematopoietic cells. Thus, the addition of FGF 98 to culture systems used for *ex vivo* expansion of cells could stimulate the rate at which certain populations of cells multiply or differentiate, and improve the effectiveness of these expansion systems in generating cells needed for transplant.

It is also believed that FGF 98 can be used for the *ex vivo* expansion of precursor cells in the nervous system. Transplant or engraftment of cells is currently being explored as a therapy for diseases in which certain populations of neurons are lost due to degeneration such as, for example, in Parkinson's disease (Bjorklund, *Curr. Opin. Neurobiol.* 2:683-689, 1992 which is incorporated by reference). Neuronal precursor cells can be

obtained from animal or human donors or from human fetal tissue and then expanded in culture using FGF 98 or other growth factors. These cells can then be engrafted into patients where they would function to replace some of the cells lost due to degeneration. Because neurotrophins have been shown to be capable of stimulating the survival and proliferation of neuronal precursors cells such as, for example, NT-3 stimulation of sympathetic neuroblast cells (Birren et al., *Develop.* 119:597-610, 1993 which is incorporated by reference), FGF 98 could also function in similar ways during the development of the nervous system and could be useful in the ex vivo expansion of neuronal cells.

In a number of circumstances it would be desirable to determine the levels of FGF 98 in a patient. The identification of FGF 98 along with the present report showing that FGF 98 is expressed by a number of tissues provides the basis for the conclusion that the presence of FGF 98 serves a normal physiologic function related to cell growth and survival. Indeed, other neurotrophic factors are known to play a role in the function of neuronal and non-neuronal tissues. (For review see Scully and Otten, *Cell Bol. Int.* 19:459-469, 1995; Otten and Gadiant, *Int. J. Devl. Neurosciences* 13:147-151, 1995 which are incorporated by reference). Endogenously produced FGF 98 may also play a role in certain disease conditions, particularly where there is cellular degeneration such as in neurodegenerative conditions or diseases. Other neurotrophic factors are known to change during disease conditions. For example, in multiple sclerosis, levels of NGF protein in the cerebrospinal fluid are increased during acute phases of the disease (Bracci-Laudiero et al., *Neuroscience Lett.* 147:9-12, 1992 which is incorporated by reference) and in systemic lupus erythematosus there is a correlation between inflammatory episodes and NGF levels in sera (Bracci-Laudiero et al. *NeuroReport* 4:563-565, 1993 which is incorporated by reference).

Given that FGF 98 is expressed in developing bone and in heart tissue, it is likely that the level of FGF 98 may be altered in a variety of conditions and that quantification of FGF 98 levels would provide clinically useful information. Furthermore, in the treatment of degenerative conditions, compositions containing FGF 98 can be administered and it would likely be desirable to achieve certain target levels of FGF 98 in sera, in cerebrospinal fluid or in any desired tissue compartment. It would, therefore, be advantageous to be able to monitor

the levels of FGF 98 in a patient. Accordingly, the present invention also provides methods for detecting the presence of FGF 98 in a sample from a patient.

The term "detection" as used herein in the context of detecting the presence of FGF 98 in a patient is intended to include the determining of the amount of FGF 98 or the ability to express an amount of FGF 98 in a patient, the distinguishing of FGF 98 from other growth factors, the estimation of prognosis in terms of probable outcome of a degenerative disease and prospect for recovery, the monitoring of the FGF 98 levels over a period of time as a measure of status of the condition, and the monitoring of FGF 98 levels for determining a preferred therapeutic regimen for the patient.

To detect the presence of FGF 98 in a patient, a sample is obtained from the patient. The sample can be a tissue biopsy sample or a sample of blood, plasma, serum, CSF or the like. FGF 98 is expressed in a variety of tissues as discussed in Example 8. Samples for detecting FGF 98 can be taken from any of these tissues. When assessing peripheral levels of FGF 98, it is preferred that the sample be a sample of blood, plasma or serum. When assessing the levels of FGF 98 in the central nervous system a preferred sample is a sample obtained from cerebrospinal fluid.

In some instances it is desirable to determine whether the FGF 98 gene is intact in the patient or in a tissue or cell line within the patient. By an intact FGF 98 gene it is meant that there are no alterations in the gene such as point mutations, deletions, insertions, chromosomal breakage, chromosomal rearrangements and the like wherein such alteration might alter production of FGF 98 or alter its biological activity, stability or the like to lead to disease processes or susceptibility to cellular degenerative conditions. Thus, in one embodiment of the present invention a method is provided for detecting and characterizing any alterations in the FGF 98 gene. The method comprises providing an oligonucleotide that contains the FGF 98 cDNA, genomic DNA or a fragment thereof or a derivative thereof. By a derivative of an oligonucleotide, it is meant that the derived oligonucleotide is substantially the same as the sequence from which it is derived in that the derived sequence has sufficient sequence complementarity to the sequence from which it is derived to hybridize to the FGF 98 gene. The derived nucleotide sequence is not necessarily physically derived from the

nucleotide sequence, but may be generated in any manner including for example, chemical synthesis or DNA replication or reverse transcription or transcription.

Typically, patient genomic DNA is isolated from a cell sample from the patient and digested with one or more restriction endonucleases such as, for example, TaqI and AluI. Using the Southern blot protocol, which is well known in the art, this assay determines whether a patient or a particular tissue in a patient has an intact FGF 98 gene or an FGF 98 gene abnormality.

Hybridization to an FGF 98 gene would involve denaturing the chromosomal DNA to obtain a single-stranded DNA; contacting the single-stranded DNA with a gene probe associated with the FGF 98 gene sequence; and identifying the hybridized DNA-probe to detect chromosomal DNA containing at least a portion of a human FGF 98 gene.

The term "probe" as used herein refers to a structure comprised of a polynucleotide that forms a hybrid structure with a target sequence, due to complementarity of probe sequence with a sequence in the target region. Oligomers suitable for use as probes may contain a minimum of about 8-12 contiguous nucleotides which are complementary to the targeted sequence and preferably a minimum of about 20.

The FGF 98 gene probes of the present invention can be DNA or RNA oligonucleotides and can be made by any method known in the art such as, for example, excision, transcription or chemical synthesis. Probes may be labeled with any detectable label known in the art such as, for example, radioactive or fluorescent labels or enzymatic marker. Labeling of the probe can be accomplished by any method known in the art such as by PCR, random priming, end labeling, nick translation or the like. One skilled in the art will also recognize that other methods not employing a labeled probe can be used to determine the hybridization. Examples of methods that can be used for detecting hybridization include Southern blotting, fluorescence in situ hybridization, and single-strand conformation polymorphism with PCR amplification.

Hybridization is typically carried out at 25° - 45° C, more preferably at 32° - 40° C and more preferably at 37° - 38° C. The time required for hybridization is from about 0.25 to about 96 hours, more preferably from about one to about 72 hours, and most preferably from about 4 to about 24 hours.

FGF 98 gene abnormalities can also be detected by using the PCR method and primers that flank or lie within the FGF 98 gene. The PCR method is well known in the art. Briefly, this method is performed using two oligonucleotide primers which are capable of hybridizing to the nucleic acid sequences flanking a target sequence that lies within an FGF 98 gene and amplifying the target sequence. The terms "oligonucleotide primer" as used herein refers to a short strand of DNA or RNA ranging in length from about 8 to about 30 bases. The upstream and downstream primers are typically from about 20 to about 30 base pairs in length and hybridize to the flanking regions for replication of the nucleotide sequence. The polymerization is catalyzed by a DNA-polymerase in the presence of deoxynucleotide triphosphates or nucleotide analogs to produce double-stranded DNA molecules. The double strands are then separated by any denaturing method including physical, chemical or enzymatic. Commonly, the method of physical denaturation is used involving heating the nucleic acid, typically to temperatures from about 80°C. to 105°C. for times ranging from about 1 to about 10 minutes. The process is repeated for the desired number of cycles.

The primers are selected to be substantially complementary to the strand of DNA being amplified. Therefore, the primers need not reflect the exact sequence of the template, but must be sufficiently complementary to selectively hybridize with the strand being amplified.

After PCR amplification, the DNA sequence comprising FGF 98 or pre-pro FGF 98 or a fragment thereof is then directly sequenced and analyzed by comparison of the sequence with the sequences disclosed herein to identify alterations which might change activity or expression levels or the like.

In another embodiment, a method for detecting FGF 98 is provided based upon an analysis of tissue expressing the FGF 98 gene. Certain tissues such as those identified below in Example 8 have been found to express the FGF 98 gene. The method comprises hybridizing a polynucleotide to mRNA from a sample of tissues that normally express the FGF 98 gene. The sample is obtained from a patient suspected of having an abnormality in the FGF 98 gene or in the FGF 98 gene of particular cells. The polynucleotide comprises SEQ ID NO:4 or a derivative thereof or a fragment thereof.

To detect the presence of mRNA encoding FGF 98 protein, a sample is obtained from a patient. The sample can be from blood or from a tissue biopsy sample. The sample may be treated to extract the nucleic acids contained therein. The resulting nucleic acid from the sample is subjected to gel electrophoresis or other size separation techniques.

The mRNA of the sample is contacted with a DNA sequence serving as a probe to form hybrid duplexes. The use of a labeled probes as discussed above allows detection of the resulting duplex.

When using the cDNA encoding FGF 98 protein or a derivative of the cDNA as a probe, high stringency conditions can be used in order to prevent false positives, that is the hybridization and apparent detection of FGF 98 nucleotide sequences when in fact an intact and functioning FGF 98 gene is not present. When using sequences derived from the FGF 98 cDNA, less stringent conditions could be used, however, this would be a less preferred approach because of the likelihood of false positives. The stringency of hybridization is determined by a number of factors during hybridization and during the washing procedure, including temperature, ionic strength, length of time and concentration of formamide. These factors are outlined in, for example, Sambrook et al. (Sambrook, et al., 1989, *supra*).

In order to increase the sensitivity of the detection in a sample of mRNA encoding the FGF 98 protein, the technique of reverse transcription/polymerization chain reaction (RT/PCR) can be used to amplify cDNA transcribed from mRNA encoding the FGF 98 protein. The method of RT/PCR is well known in the art (see example 9 and Figure 6 below).

The RT/PCR method can be performed as follows. Total cellular RNA is isolated by, for example, the standard guanidium isothiocyanate method and the total RNA is reverse transcribed. The reverse transcription method involves synthesis of DNA on a template of RNA using a reverse transcriptase enzyme and a 3' end primer. Typically, the primer contains an oligo(dT) sequence. The cDNA thus produced is then amplified using the PCR method and FGF 98 specific primers. (Belyavsky et al., *Nucl. Acid Res.* 17:2919-2932, 1989; Krug and Berger, *Methods in Enzymology*, 152:316-325, Academic Press, NY, 1987 which are incorporated by reference).

The polymerase chain reaction method is performed as described above using two oligonucleotide primers that are substantially complementary to the two flanking regions of the DNA segment to be amplified.

Following amplification, the PCR product is then electrophoresed and detected by ethidium bromide staining or by phosphoimaging.

The present invention further provides for methods to detect the presence of the FGF 98 protein in a sample obtained from a patient. Any method known in the art for detecting proteins can be used. Such methods include, but are not limited to immunodiffusion, immunoelectrophoresis, immunochemical methods, binder-ligand assays, immunohistochemical techniques, agglutination and complement assays. (for example, see *Basic and Clinical Immunology*, 217-262, Sites and Terr, eds., Appleton & Lange, Norwalk, CT, 1991 which is incorporated by reference). Preferred are binder-ligand immunoassay methods including reacting antibodies with an epitope or epitopes of the FGF 98 protein and competitively displacing a labeled FGF 98 protein or derivative thereof. A preferred antibody is prepared according to Example 9.

As used herein, a derivative of the FGF 98 protein is intended to include a polypeptide in which certain amino acids have been deleted or replaced or changed to modified or unusual amino acids wherein the FGF 98 derivative is biologically equivalent to FGF 98 and wherein the polypeptide derivative cross-reacts with antibodies raised against the FGF 98 protein. By cross-reaction it is meant that an antibody reacts with an antigen other than the one that induced its formation.

Numerous competitive and non-competitive protein binding immunoassays are well known in the art. Antibodies employed in such assays may be unlabeled, for example as used in agglutination tests, or labeled for use in a wide variety of assay methods. Labels that can be used include radionuclides, enzymes, fluorescers, chemilumescers, enzyme substrates or co-factors, enzyme inhibitors, particles, dyes and the like for use in radioimmunoassay (RIA), enzyme immunoassays, *e.g.*, enzyme-linked immunosorbent assay (ELISA), fluorescent immunoassays and the like.

Polyclonal or monoclonal antibodies to the FGF 98 protein or an epitope thereof can be made for use in immunoassays by any of a number of methods known in the art.

By epitope reference is made to an antigenic determinant of a polypeptide. An epitope could comprise 3 amino acids in a spacial conformation which is unique to the epitope. Generally an epitope consists of at least 5 such amino acids. Methods of determining the spatial conformation of amino acids are known in the art, and include, for example, x-ray crystallography and 2 dimensional nuclear magnetic resonance.

One approach for preparing antibodies to a protein is the selection and preparation of an amino acid sequence of all or part of the protein, chemically synthesizing the sequence and injecting it into an appropriate animal, usually a rabbit or a mouse (See Example 9).

Oligopeptides can be selected as candidates for the production of an antibody to the FGF 98 protein based upon the oligopeptides lying in hydrophilic regions, which are thus likely to be exposed in the mature protein. A preferred oligopeptide is KRYPKGQPELQKPFK (SEQ ID NO:6).

Antibodies to FGF 98 can also be raised against oligopeptides that include one or more of the conserved regions identified herein such that the antibody can cross-react with other family members. Such antibodies can be used to identify and isolate the other family members.

Methods for preparation of the FGF 98 protein or an epitope thereof include, but are not limited to chemical synthesis, recombinant DNA techniques or isolation from biological samples. Chemical synthesis of a peptide can be performed, for example, by the classical Merrifield method of solid phase peptide synthesis (Merrifield, *J. Am. Chem. Soc.* 85:2149, 1963 which is incorporated by reference) or the FMOC strategy on a Rapid Automated Multiple Peptide Synthesis system (E. I. du Pont de Nemours Company, Wilmington, DE) (Caprino and Han, *J Org Chem* 37:3404, 1972 which is incorporated by reference).

Polyclonal antibodies can be prepared by immunizing rabbits or other animals by injecting antigen followed by subsequent boosts at appropriate intervals. The animals are bled and sera assayed against purified FGF 98 protein usually by ELISA or by bioassay based upon the ability to block the action of FGF 98 on neurons or other cells. When using avian species, e.g., chicken, turkey and the like, the antibody can be isolated from the yolk of the

egg. Monoclonal antibodies can be prepared after the method of Milstein and Kohler by fusing splenocytes from immunized mice with continuously replicating tumor cells such as myeloma or lymphoma cells. (Milstein and Kohler, *Nature* 256:495-497, 1975; Galfre and Milstein, *Methods in Enzymology: Immunochemical Techniques* 73:1-46, Langone and Banatis eds., Academic Press, 1981 which are incorporated by reference). The hybridoma cells so formed are then cloned by limiting dilution methods and supernates assayed for antibody production by ELISA, RIA or bioassay.

The unique ability of antibodies to recognize and specifically bind to target proteins provides an approach for treating an over expression of the protein. Thus, another aspect of the present invention provides for a method for preventing or treating diseases involving over expression of the FGF 98 protein by treatment of a patient with specific antibodies to the FGF 98 protein.

Specific antibodies, either polyclonal or monoclonal, to the FGF 98 protein can be produced by any suitable method known in the art as discussed above. For example, murine or human monoclonal antibodies can be produced by hybridoma technology or, alternatively, the FGF 98 protein, or an immunologically active fragment thereof, or an anti-idiotypic antibody, or fragment thereof can be administered to an animal to elicit the production of antibodies capable of recognizing and binding to the FGF 98 protein. Such antibodies can be from any class of antibodies including, but not limited to IgG, IgA, IgM, IgD, and IgE or in the case of avian species, IgY and from any subclass of antibodies.

Polypeptides encoded by the instant polynucleotides and corresponding full-length genes can be used to screen peptide libraries, protein libraries, small molecule libraries, and phage display libraries, and other known methods, to identify analogs or antagonists.

Native FGF polypeptides may play a role in cancer. For example, FGF family members can induce marked morphological transformation of NIH 3T3 cells, and exhibit strong tumorigenicity in nude mice. Angiogenic activity has been exhibited by FGF family members. Thus, inhibitors of FGF can be used to treat cancer, such as prostate.

A library of peptides may be synthesized following the methods disclosed in U.S. Patent No. 5,010,175, and in PCT No. WO 91/17823. As described below in brief, a mixture of peptides is prepared, which is then screened to identify the peptides exhibiting the

desired signal transduction and receptor binding activity. According to the method of the '175 patent, a suitable peptide synthesis support (*e.g.*, a resin) is coupled to a mixture of appropriately protected, activated amino acids. The concentration of each amino acid in the reaction mixture is balanced or adjusted in inverse proportion to its coupling reaction rate so that the product is an equimolar mixture of amino acids coupled to the starting resin. The bound amino acids are then deprotected, and reacted with another balanced amino acid mixture to form an equimolar mixture of all possible dipeptides. This process is repeated until a mixture of peptides of the desired length (*e.g.*, hexamers) is formed. Note that one need not include all amino acids in each step: one may include only one or two amino acids in some steps (*e.g.*, where it is known that a particular amino acid is essential in a given position), thus reducing the complexity of the mixture. After the synthesis of the peptide library is completed, the mixture of peptides is screened for binding to the selected polypeptide. The peptides are then tested for their ability to inhibit or enhance activity. Peptides exhibiting the desired activity are then isolated and sequenced.

The method described in PCT No. WO 91/17823 is similar. However, instead of reacting the synthesis resin with a mixture of activated amino acids, the resin is divided into twenty equal portions (or into a number of portions corresponding to the number of different amino acids to be added in that step), and each amino acid is coupled individually to its portion of resin. The resin portions are then combined, mixed, and again divided into a number of equal portions for reaction with the second amino acid. In this manner, each reaction may be easily driven to completion. Additionally, one may maintain separate "subpools" by treating portions in parallel, rather than combining all resins at each step. This simplifies the process of determining which peptides are responsible for any observed receptor binding or signal transduction activity.

In such cases, the subpools containing, *e.g.*, 1-2,000 candidates each are exposed to one or more polypeptides of the invention. Each subpool that produces a positive result is then resynthesized as a group of smaller subpools (sub-subpools) containing, *e.g.*, 20-100 candidates, and reassayed. Positive sub-subpools may be resynthesized as individual compounds, and assayed finally to determine the peptides that exhibit a high binding constant. These peptides can be tested for their ability to inhibit or enhance the native activity. The

methods described in PCT No. WO 91/7823 and U.S. Patent No. 5,194,392 (herein incorporated by reference) enable the preparation of such pools and subpools by automated techniques in parallel, such that all synthesis and resynthesis may be performed in a matter of days.

Peptide agonists or antagonists are screened using any available method, such as signal transduction, antibody binding, receptor binding and mitogenic assays. The assay conditions ideally should resemble the conditions under which the native activity is exhibited *in vivo*, that is, under physiologic pH, temperature, and ionic strength. Suitable agonists or antagonists will exhibit strong inhibition or enhancement of the native activity at concentrations that do not cause toxic side effects in the subject. Agonists or antagonists that compete for binding to the native polypeptide may require concentrations equal to or greater than the native concentration, while inhibitors capable of binding irreversibly to the polypeptide may be added in concentrations on the order of the native concentration.

The end results of such screening and experimentation will be at least one novel polypeptide binding partner, such as a receptor, encoded by a gene or a cDNA of the invention, and at least one peptide agonist or antagonist of the novel binding partner. Such agonists and antagonists can be used to modulate, enhance, or inhibit receptor function in cells to which the receptor is native, or in cells that possess the receptor as a result of genetic engineering. Further, if the novel receptor shares biologically important characteristics with a known receptor, information about agonist/antagonist binding may help in developing improved agonists/antagonists of the known receptor.

The therapeutic FGF 98 polynucleotides and polypeptides of the present invention may be utilized in gene delivery vehicles. The gene delivery vehicle may be of viral or non-viral origin (see generally, Jolly, *Cancer Gene Therapy* 1:51-64 (1994); Kimura, *Human Gene Therapy* 5:845-852 (1994); Connelly, *Human Gene Therapy* 1:185-193 (1995); and Kaplitt, *Nature Genetics* 6:148-153 (1994)). Gene therapy vehicles for delivery of constructs including a coding sequence of a therapeutic of the invention can be administered either locally or systemically. These constructs can utilize viral or non-viral vector approaches. Expression of such coding sequences can be induced using endogenous

mammalian or heterologous promoters. Expression of the coding sequence can be either constitutive or regulated.

The present invention can employ recombinant retroviruses which are constructed to carry or express a selected nucleic acid molecule of interest. Retrovirus vectors that can be employed include those described in EP 0 415 731; WO 90/07936; WO 94/03622; WO 93/25698; WO 93/25234; U.S. Patent No. 5,219,740; WO 93/11230; WO 93/10218; Vile and Hart, *Cancer Res.* 53:3860-3864 (1993); Vile and Hart, *Cancer Res.* 53:962-967 (1993); Ram et al., *Cancer Res.* 53:83-88 (1993); Takamiya et al., *J. Neurosci. Res.* 33:493-503 (1992); Baba et al., *J. Neurosurg.* 79:729-735 (1993); U.S. Patent No. 4,777,127; GB Patent No. 2,200,651; and EP 0 345 242. Preferred recombinant retroviruses include those described in WO 91/02805.

Packaging cell lines suitable for use with the above-described retroviral vector constructs may be readily prepared (see PCT publications WO 95/30763 and WO 92/05266), and used to create producer cell lines (also termed vector cell lines) for the production of recombinant vector particles. Within particularly preferred embodiments of the invention, packaging cell lines are made from human (such as HT1080 cells) or mink parent cell lines, thereby allowing production of recombinant retroviruses that can survive inactivation in human serum.

The present invention also employs alphavirus-based vectors that can function as gene delivery vehicles. Such vectors can be constructed from a wide variety of alphaviruses, including, for example, Sindbis virus vectors, Semliki forest virus (ATCC VR-67; ATCC VR-1247), Ross River virus (ATCC VR-373; ATCC VR-1246) and Venezuelan equine encephalitis virus (ATCC VR-923; ATCC VR-1250; ATCC VR 1249; ATCC VR-532). Representative examples of such vector systems include those described in U.S. Patent Nos. 5,091,309; 5,217,879; and 5,185,440; and PCT Publication Nos. WO 92/10578; WO 94/21792; WO 95/27069; WO 95/27044; and WO 95/07994.

Gene delivery vehicles of the present invention can also employ parvovirus such as adeno-associated virus (AAV) vectors. Representative examples include the AAV vectors disclosed by Srivastava in WO 93/09239, Samulski et al., *J. Vir.* 63:3822-3828

(1989); Mendelson et al., *Virol.* 166:154-165 (1988); and Flotte et al., *P.N.A.S.* 90:10613-10617 (1993).

Representative examples of adenoviral vectors include those described by Berkner, *Biotechniques* 6:616-627 (Biotechniques); Rosenfeld et al., *Science* 252:431-434 (1991); WO 93/19191; Kolls et al., *P.N.A.S.* :215-219 (1994); Kass-Eisler et al., *P.N.A.S.* 90:11498-11502 (1993); Guzman et al., *Circulation* 88:2838-2848 (1993); Guzman et al., *Cir. Res.* 73:1202-1207 (1993); Zabner et al., *Cell* 75:207-216 (1993); Li et al., *Hum. Gene Ther.* 4:403-409 (1993); Cailaud et al., *Eur. J. Neurosci.* 5:1287-1291 (1993); Vincent et al., *Nat. Genet.* 5:130-134 (1993); Jaffe et al., *Nat. Genet.* 1:372-378 (1992); and Levrero et al., *Gene* 101:195-202 (1992). Exemplary adenoviral gene therapy vectors employable in this invention also include those described in WO 94/12649, WO 93/03769; WO 93/19191; WO 94/28938; WO 95/11984 and WO 95/00655. Administration of DNA linked to killed adenovirus as described in Curiel, *Hum. Gene Ther.* 3:147-154 (1992) may be employed.

Other gene delivery vehicles and methods may be employed, including polycationic condensed DNA linked or unlinked to killed adenovirus alone, for example Curiel, *Hum. Gene Ther.* 3:147-154 (1992); ligand-linked DNA, for example see Wu, *J. Biol. Chem.* 264:16985-16987 (1989); eukaryotic cell delivery vehicles cells, for example see U.S. Serial No. 08/240,030, filed May 9, 1994, and U.S. Serial No. 08/404,796; deposition of photopolymerized hydrogel materials; hand-held gene transfer particle gun, as described in U.S. Patent No. 5,149,655; ionizing radiation as described in U.S. Patent No. 5,206,152 and in WO 92/11033; nucleic charge neutralization or fusion with cell membranes. Additional approaches are described in Philip, *Mol. Cell Biol.* 14:2411-2418 (1994), and in Woffendin, *Proc. Natl. Acad. Sci.* 91:1581-1585 (1994).

Naked DNA may also be employed. Exemplary naked DNA introduction methods are described in WO 90/11092 and U.S. Patent No. 5,580,859. Uptake efficiency may be improved using biodegradable latex beads. DNA coated latex beads are efficiently transported into cells after endocytosis initiation by the beads. The method may be improved further by treatment of the beads to increase hydrophobicity and thereby facilitate disruption of the endosome and release of the DNA into the cytoplasm. Liposomes that can act as gene

delivery vehicles are described in U.S. Patent No. 5,422,120, PCT Patent Publication Nos. WO 95/13796, WO 94/23697, and WO 91/14445, and EP No. 0 524 968.

Further non-viral delivery suitable for use includes mechanical delivery systems such as the approach described in Woffendin et al., *Proc. Natl. Acad. Sci. USA* 91(24):11581-11585 (1994). Moreover, the coding sequence and the product of expression of such can be delivered through deposition of photopolymerized hydrogel materials. Other conventional methods for gene delivery that can be used for delivery of the coding sequence include, for example, use of hand-held gene transfer particle gun, as described in U.S. Patent No. 5,149,655; use of ionizing radiation for activating transferred gene, as described in U.S. Patent No. 5,206,152 and PCT Patent Publication No. WO 92/11033.

FGF has been implicated in diseases characterized by loss of function, inadequate function/number, abnormal function or death of cells, tissues or organs for which function or survival can be prolonged/rescued, and abnormalities reversed or prevented by therapy with FGF.

According to the invention, FGF 98 is likely to play a role in degenerative and other neurological diseases (disease characterized by loss of function or death of CNS cells, including neurons, glial cells, astrocytes, and others), in which therapy may prolong survival of cells or rescue/preserve function. Examples include Parkinson's disease, Alzheimers or other dementia, stroke, trauma, sequelae of infection, epilepsy, cranial nerve injury or dysfunction. FGF 98 may also be useful in treatment of eye disease characterized by similar responses of retinal cells, structural cells, nerve cells, etc., to administration of the molecule. Examples include Retinitis Pigmentosa, age-related macular degeneration, retinal detachment, ischemic retinal disease, etc.

FGF 98 may be useful in treatment of disorders of hearing, smell or taste due to loss/dysfunction of cells responsive to the molecule, including cranial or peripheral nerves, sensory cells, supporting cells, etc., and in treatment of peripheral motor or sensor nerve disease characterized by loss of function or death of neurons, nerves, other CNS or PNS cells or supporting cells: Amyotrophic Lateral Sclerosis, motor or Sensory Neuropathies (due to many causes; viral infections, metabolic disease-diabetes, autoimmune insult, idiopathic, degenerative, etc.) trauma.

Ischemic vascular disease may be amenable to FGF 98 treatment, wherein the disease is characterized by inadequate blood flow to an organ(s). Treatment may induce therapeutic angiogenesis or preserve function/survival of cells (myocardial ischemia/infarction, peripheral vascular disease, renal artery disease, stroke). Cardiomyopathies characterized by loss of function or death of cardiac myocytes or supporting cells in the heart (congestive heart failure, myocarditis) may also be treated using FGF 98, as can musculoskeletal disease characterized by loss of function, inadequate function or death of skeletal muscle cells, bone cells or supporting cells. Examples include skeletal myopathies, bone disease, and arthritis.

FGF 98 polynucleotides and polypeptides may aid in correction of congenital defects due to loss of FGF 98 molecule or its function (heart, lung, brain, limbs, kidney, etc.).

Treatment of wound healing is yet another use of FGF 98 polypeptides and polynucleotides, either due to trauma, disease, medical or surgical treatment, including regeneration of cell populations and tissues depleted by these processes. Examples include liver regeneration, operative wound healing, re-endothelialization of injured blood vessels, healing of traumatic wounds, healing of ulcers due to vascular, metabolic disease, etc., bone fractures, loss of cells due to inflammatory disease, etc.

FGF 98 may also be used in screens to identify drugs for treatment of cancers which involve over activity of the molecule, or new targets which would be useful in the identification of new drugs.

For all of the preceding embodiments, the clinician will determine, based on the specific condition, whether FGF 98 polypeptides or polynucleotides, antibodies to FGF 98, or small molecules such as peptide analogues or antagonists, will be the most suitable form of treatment. These forms are all within the scope of the invention.

Preferred embodiments of the invention are described in the following examples. Other embodiments within the scope of the claims herein will be apparent to one skilled in the art from consideration of the specification or practice of the invention as disclosed herein. It is intended that the specification, together with the examples, be considered exemplary only, with the scope and spirit of the invention being indicated by the claims which follow the examples.

EXAMPLES

EXAMPLE 1

EXPRESSION OF FGF 98 IN INSECT CELLS

The baculovirus vectors for expression of FGF 98 were generated by PCR of the coding region from a Bluescript vector containing the cDNA for FGF 98, prepared from a human heart cDNA library. Restriction sites for PstI and KpnI were added to facilitate directional subcloning into Bluebac 4.5 baculovirus vector (Invitrogen).

Supernatants from insect cells were generated for purification and bioassay by infecting SF9 cells at approximately $1-2 \times 10^6$ /ml with recombinant baculovirus derived from these plasmids. A multiplicity of infection (moi) of 2-5 was used for the 48 hour infections. SF9 cells were removed from the supernatant by centrifugation and the supernatants further clarified by $0.2 \mu\text{M}$ sterile filtration. In order to determine high producers, SF9 cells infected with different clones of baculovirus supernatants were analyzed by Western blot (Figure 1). A peptide antibody directed against the C terminal region of FGF 98 (AA177 to 195) was used. The yield in insect cells was 4 to 8 mg protein per liter of CM.

EXAMPLE 2

ISOLATION OF FGF 98

Recombinant FGF 98 from the conditioned medium of SF-9 cells was purified by Heparin Sepharose (HS) affinity chromatography, followed by Mono S cation exchange chromatography. The recombinant FGF 98 bound to the HS was eluted using a step-wise salt gradient. Next, the recombinant FGF 98 eluted from the HS column was bound to the Mono S cation exchange column and was eluted using a nonlinear salt gradient.

Further details of the procedure utilized are described below.

A. Heparin Sepharose® Affinity Chromatography

The conditioned medium from Example 1 (30 ml) was loaded through a 5 ml bed of HS resin. The column was equilibrated in a buffer containing 150 mM NaCl and 10

mM Tris-HCl at pH 7.3. Once the conditioned medium was loaded, the column was washed extensively with the equilibration buffer until the absorbency at 280 nm returned to baseline. The protein was eluted from the HS column with an increasing step-wise NaCl gradient. The NaCl concentrations were 0.5M, 1M, and 2M NaCl, in 10 mM Tris-HCl at pH 7.3. The flow rate of the column during elution was approximately 30 ml/hr and 0.5-ml size fractions were collected.

The fractions were tested for FGF 98 bioactivity utilizing granulosa cells. The assay is described in Section 4 below. The fractions with the highest bioactivity were eluted with 2M NaCl (Figure 2).

When analyzed by SDS-PAGE followed by Coomassie blue staining the peak 2M NaCl fractions gave a single band with a MW of 28 kDa (Figure 3A). When analyzed by Western using an anti-FGF 98 peptide antibody the 2M NaCl fractions were immunoactive with a single band with a MW of 28 kDa (Figure 3B). No immunoreactive bands could be observed with the 1M NaCl fractions.

The homogeneity of the pooled 2M NaCl fractions was also analyzed by RP-HPLC. A single major protein peak eluting at 40% acetonitrile was observed. Closely associated with it was a minor protein peak that accounted for less than 5% of the protein mass (Figure 4). When the HPLC fractions were analyzed for bioactivity using granuloma cells, both protein peaks were active. When analyzed by SDS-PAGE followed by Coomassie blue staining, both fractions gave a single band with an apparent MW of 28k (Figure 5A). When this was followed by Western analysis both bands were recognized by a peptide antibody specific for FGF 98 (Figure 5B).

However, when the bioactivity of the pooled 2M NaCl fractions was analyzed on vascular endothelial cells, cytotoxicity was observed, indicating the presence of endotoxin-like substances. Those were eliminated using Mono S cation exchange chromatography as described below.

B. Large scale isolation of FGF 98.

A 3-liter batch of SF-9 cell-conditioned medium with pH adjusted to 7.3 and NaCl concentration raised to 0.5M was then allowed to run overnight at 4°C through a 30 ml

bed of HS resin. The column was equilibrated in a buffer containing 0.5M NaCl and 10 mM Tris- HCl pH 7.3. Following loading, the column was washed with the same buffer. The bound proteins were then eluted with 1M and 2M NaCl in 10 mM Tris pH 7.3. The flow rate of the column during the elution was 90 ml/hr and 3 ml size reactions were collected (Figures 6A, B).

When the collected fractions were tested for bioactivity using granulosa cells, all of the bioactivity was present in the 2M NaCl fractions. When the 1M and 2M NaCl fractions were analyzed by SDS-PAGE, the 1M NaCl fraction showed multiple bands with MW ranging from 200 kDa to 8 kDa. In contrast, the 2M NaCl fractions show a major single band with a MW of 28 kDa (Figure 7A). When analyzed by Western using an FGF 98 peptide antibody, the 1M NaCl contained a very weak immunoreactive band with a MW of 28 kDa. In contrast the 2M NaCl showed a very strong immunoreactive band at 28 kDa together with a weak band at 56 kDa (Figure 7B).

When the 2M NaCl fractions were tested on vascular endothelial cells, all fractions were toxic, indicating the presence of endotoxin-like material. Those were eliminated using Mono S cation exchange chromatography (Figure 8).

C. Mono S Cation Exchange Chromatography

The pooled 2M NaCl fractions eluted from the HS column were diluted ten-fold with 10 mM Tris-HCl at pH 7.3 to a final salt concentration of 0.2M NaCl. The material was then loaded with a Super loop onto a Mono S column linked to a FPLC system (Pharmacia, Piscataway, NJ). The Mono S cation exchange column was equilibrated with 10 mM Tris-HCl at pH 7.2. When the pooled fractions were loaded, the column was washed extensively at a flow rate of 1 ml/min. with the equilibration buffer until the absorbency returned to baseline. Then, protein was eluted from the column with a nonlinear NaCl gradient, 0.2M to 2M NaCl in 10 mM Tris-HCl at pH 7.3 at a flow rate of 1 ml/min. and 1 ml fractions were collected.

One major protein peak of activity was observed that eluted at about 0.8M NaCl (Figure 8). Fractions across the protein peak were assayed for bioactivity and subjected

to SDS-PAGE analysis. The results of the gel analysis (not illustrated) showed that protein from NaCl fractions exhibited an apparent molecular weight of 28 kD.

Western analysis of the Mono S 0.8M pooled fraction with peptide antibodies directed against AA176 to AA190 of FGF 98 revealed a single band with an apparent MW of 28 kD.

The homogeneity of the pooled Mono S fractions was also analyzed by RP-HPLC using a C4 column equilibrated in 0.1% TFA, 20% acetonitrile, using an acetonitrile gradient of 20 to 100%. A profile similar to that shown in Figure 4 was obtained (data not shown).

EXAMPLE 3

FGF 98 BIOACTIVITY EFFECT ON GRANULOSA CELLS

Stock cultures of granulosa cells were grown and maintained in Dulbecco's modified Eagle medium supplemented with 10% bovine serum, 0.25 μ l/ml fungizone, and 10 ng/ml FGF. The cells were incubated at 37°C in a 10% CO₂ atmosphere with 99% humidity. For the bioactivity assay, the cells were seeded in 12-well plates at a density of 5×10^3 cells per well in 1 ml of medium as described above for the stock cultures, and as described in Gospodarowicz et al. *Meth. Enzymol.* 147B:106-119 (D. Barnes et al., ed.) Academic Press, Orlando, FL (1987).

Ten microliter aliquots of the desired column fractions were diluted into 1 ml of 0.2% (w/v) gelatin in phosphate buffered saline (PBS). Ten microliters of this dilution were added to granulosa cells seeded in 12-well cluster plates containing 22 mm wells, at 5×10^3 cells per well, and a 10 μ l aliquot of either the diluted column fractions or medium containing 10 ng bFGF were added to the cells every other day.

After five days in culture, the cells were trypsinized and the final cell density was determined using a Coulter® counter. (Coulter Electronics, Hialeah, FL, USA). The cells were released from the plates by replacing the culture medium with a solution containing 0.9% NaCl, 0.01M sodium phosphate (pH 7.4), 0.05% trypsin, and 0.02% EDTA (STV). The cells were incubated in this solution for 5-10 minutes at 37°C, and then the stock culture medium was added to the cells. The cells were then counted using a Coulter® counter.

Results shown in Figure 9 demonstrates the effect of different dilutions of FGF 98 on granulosa cells as compared to basic FGF. The figure legend for Figure 9 is as follows: O=basic FGF, =FGF 98.

The final cell density was graphed as a function of protein concentration. The protein concentration is graphed on a log scale. The protein concentration was determined by Bradford assay according to instructions accompanying the protein assay kit from BIORAD (Richmond, CA, USA).

The ED_{50} was calculated by (a), dividing in half the difference between the lowest and highest cell density value of the curve; and (b), determining from the graph what protein concentration corresponds to that cell density number obtained in (a). According to Figure 9 the ED_{50} for bFGF was found to be approximately 40 pg/ml, that for FGF 98 was about 100 ng/ml. The results of this assay demonstrate that FGF 98 was about 2500-fold less active than bFGF.

EXAMPLE 4

ABSENCE OF FGF 98 BIOACTIVITY ON ABAE OR ACE CELLS

FGF 98 can be characterized by its lack of activity on vascular endothelial cells derived from large vessels ("ABAE") or capillary cells ("ACE") as compared with basic FGF ("bFGF"). Stock cultures of ABAE and ACE cells were grown and maintained in Dulbecco's modified Eagle medium supplemented with 10% bovine serum, 0.25 μ l/ml fungizone, and 2 ng/ml bFGF. The cells were incubated at 37°C with a 10% CO₂ concentration and 99% humidity.

In the mitogenic assay, either 10⁴ ABAE or 5x10³ ACE cells were plated per well in 12-well plates in stock culture medium, as described in Gospodarowicz et al., *Proc. Natl. Acad. USA* 73:4120-4124 (1976), Gospodarowicz et al., *J. Cell. Physiol.* 127:121-136 (1976); and Gospodarowicz et al., *Proc. Natl. Acad. USA* 86:7311-7315 (1989). Ten microliter aliquots of the fractions to be tested were diluted into 1 ml of 0.2% (w/v) gelatin in phosphate buffered saline. Ten microliters of this dilution were added to cells seeded in 12-well cluster plates. A 10 μ l aliquot of either the diluted purified FGF 98 medium-containing fractions or 2 ng bFGF was added every other day.

After five days in culture, the cells were trypsinized and the final cell density was determined using a Coulter® counter. The cells were released from the plates by replacing the medium with a solution containing 0.9% NaCl, 0.01%M sodium phosphate (pH 7.4), 0.05% trypsin, and 0.02% EDTA (STV). The cells were incubated in this solution for 5-10 minutes at 37°C, and then the stock culture medium was added to the cells. The cells then were counted using a Coulter® counter (Coulter Electronics, Hialeah, FL, USA).

In the absence of heparin, results demonstrated the lack of activity of FGF 98 on ABAE and ACE cells in contrast to bFGF. In the presence of heparin (2 µg/ml), FGF 98 was bioactive on ABAE as well as ACE cells.

EXAMPLE 5

FGF 98 EXPRESSION PATTERNS IN EMBRYOGENESIS

To determine the expression pattern of FGF 98 during embryogenesis, wholemount in situ hybridizations were performed on mouse embryos ranging in stage from embryonic day (E) 7.5 to E11.5 and on limbs from E12.5 to E14.5. For wholemount RNA in situ hybridizations, mouse embryos (CD-1) were isolated, fixed, and processed following the protocol by Nieto, M.A., Patel, K., Wilkinson, D.G. 1995. In situ hybridization analysis of chick embryos in whole mount and tissue section. In: *Methods in Avian Embryology* (ed. M. Bronner-Fraser), San Diego, Academic Press. Antisense riboprobe for FGF 98 was synthesized using the cDNA corresponding to EST accession number AA656693 as a template, linearized with XbaI and transcribed with T7 RNA polymerase. This probe encompasses C-terminal coding sequence and 3' untranslated region. The *fgf-8* probe was prepared as described by Crossley et al., *Development* 121:439-451 (1995). Embryos processed for wholemount in situ hybridization were equilibrated in 30% sucrose, embedded in OCT or 15% sucrose, 7.5% gelatin and cryosectioned at 12 or 30 µm, respectively.

At E7.5, no FGF 98 transcripts were detectable. At E8.5, FGF 98 transcripts were present in the paraxial mesoderm and in the newly formed somites. At all subsequent stages analyzed, FGF 98 continued to be expressed in the rostral part of the paraxial mesoderm and throughout the newly formed somites. As the more rostral somites mature,

FGF 98 transcripts first become restricted to the caudal part of the somite, and then to the caudal part of the myotome.

In the developing limbs, FGF 98 expression was initiated at E10.5 in a dorsal and a ventral domain in the mesenchyme. At E11, two separate domains of expression were observed on the ventral side of the limb. From E11.5 to at least E14.5, FGF 98 expression appeared to outline the differentiating skeletal elements, first the humerus, radius and ulna, followed by the digits. Transverse sections through E13.5 digits demonstrated that FGF 98 expression was present in tissues surrounding the differentiating cartilage elements. Furthermore, FGF was expressed in the interdigital regions at E12.5.

In the developing nervous system, FGF 98 transcripts were detected from E9.0 through at least E11.5. From E9.0 to E9.5, FGF 98 was transiently expressed in the caudal part of the neural plate which has not yet closed. In the differentiating neural tube, FGF 98 expression was initiated at E 9.5, in a dorsal and a ventral domain in the mantle layer. Furthermore, FGF 98 expression was observed in the dorsal root ganglia and in a subset of cranial ganglia (v, vii-viii, ix-x).

During the early development of the brain, the expression pattern of FGF 98 resembled that of *fgf-8*. At E8.5, FGF 98 expression was restricted to a narrow stripe in the head similar to a domain observed for *fgf-8* transcripts (Crossley et al., *Development* 121:439-451 (1995)). At subsequent stages, this FGF 98 expression domain marked the border between the midbrain and the hindbrain. However, the expression domain of FGF 98 in this region appeared to be wider than that of *fgf-8*. In the commissural plate of the telencephalon, FGF 98 expression was initiated at E9.0, approximately 0.5 days later than that of *fgf-8*. In the diencephalon, FGF 98 transcripts were first detected at E9.5 in both a dorsal and a ventral domain, and by E10.5, dorsal expression extended from the diencephalon rostrally into the telencephalon. In the optic stalks, FGF 98 expression was restricted ventrally, similar to *fgf-8*.

Finally, FGF 98 mRNA was detected in the pharyngeal region beginning at E9.5 in a restricted region of the second and the third branchial arch.

EXAMPLE 6

STEM CELL PROLIFERATION ASSAY

The neuron stem cell/progenitor assay involved the culturing of hippocampal cells from the embryonic rat brain for 6 days at 37°C in medium containing FGF 98, basic FGF, or no added factor. Neuronal survival and proliferation were assessed visually under a microscope on day 6. Photomicrographs were made and the cell number in each condition counted (the data are shown in Figure 10; series 1, 6 days in vitro DIV). The cultures were then washed and fresh media added without added FGF 98 or bFGF. Following 15 days in media without added factors cells were visually inspected, photomicrographs made and cells counted (Figure 11, series 2, 21 DIV). FGF 98 increases the number of neural cells at both 6DIV and 21DIV. FGF 98 stimulates the proliferation of neural progenitors under these conditions. The effects of FGF 98 are significantly different than the effect of bFGF where there is proliferation at 6 DIV but many of these cells die following withdraw of the growth factor. Few cells die in the FGF 98 treatment conditions.

Based on these and other results, FGF 98 can be used for the *ex vivo* expansion of precursor cells on the nervous system. Transplant or engraftment of cells can be used as a therapy for diseases in which specific populations of neurons are lost due to degeneration such as, for example, Parkinson's disease and Alzheimer's disease.

EXAMPLE 7

PREPARATION OF PRIMARY HIPPOCAMPAL CELL CULTURES

Hippocampi were dissected from E18-19 rats. The tissue was collected in Hibernate™ media (Gibco BRL) and minced into approximately 1mm pieces. Tissue was dissociated using the Papain Dissociation System (Worthington Biochemical Corporation). Following isolation the cells were resuspended in serum-free media composed of Neurobasal™ (Gibco BRL), 2%B27 supplement (GibcoBRL), L-glutamate and antibiotics. At the time of plating the growth factors were added: bFGF at 10ng/ml or FGF 98 at 200ng/ml. Cells were plated in 35mm tissue culture dishes coated with poly-L-lysine at a concentration of 7.5×10^4 cells per dish. Following 6 days at 37°C in 5%CO₂ cells were rinsed

and fed with fresh media; no growth factors were added. Fifteen days after the factors were removed (total 21 DIV) the cells were fixed for 30 min in 4% paraformaldehyde and photographed. Some cultures are stained with the monoclonal antibody to the nestin intermediate filament (Developmental Studies Hybridoma Bank). This antibody recognizes neural stem cells/precursors. (Johe, K.K., et al. (1996). *Genes and Development* 10:3129-3140.) Figure 11 shows the expression of nestin in 15 DIV cultures treated with either bFGF or FGF 98 for the first 6 days. There are many more nestin positive cells in the FGF 98 treated cultures. See Figure 11, Panel A, nestin expression in bFGF treated cultures, and Figure 11, Panel B, nestin expression in FGF 98 treated cultures.

EXAMPLE 8

FGF 98 EXPRESSION

A survey of FGF 98 expression was performed in mouse embryonic and adult tissues (>3months) using in situ hybridization. FGF 98 is expressed in the E14 mouse nervous system; areas of expression include the motor neurons of the brainstem and spinal cord, the dorsal root and cranial ganglia of the peripheral nervous system, the ventral tegmentum, developing forebrain, the region of the isthmus and developing cerebellum. In the adult brain FGF 98 is expressed in both the brain and spinal cord including the motor neurons, the substantia nigra, the deep cerebellar nuclei, the red nucleus and other areas. There are also several non-neuronal tissues in which FGF 98 is expressed including lung, heart, developing bone and chondrocytes.

Figure 12, Panel A, shows *in situ* hybridization of the adult brain stem and cerebellum. FGF 98 mRNA expression in deep cerebellar nuclei, and motor neurons of the facial and cochlear nuclei is visible. Figure 12, Panel B shows a high magnification of FGF 98 expression in motor neurons of the facial nucleus.

EXAMPLE 9

PREPARATION OF ANTISERA TO FGF 98 BY IMMUNIZATION OF RABBITS
WITH AN FGF 98 PEPTIDE.

The peptide sequence corresponding to amino acids 176-190 of the human FGF 98 protein was synthesized and coupled to keyhole limpet hemocyanin (KLH) as described (Harlow and Land, Antibodies: A Laboratory Manual, 1988. Cold Spring Harbor Laboratory, New York, NY) The KLH-coupled peptide was submitted to EL Labs and each of three rabbits were immunized. (EL Labs, Soquel, CA).

Peptide sequence: H-KRYPKGQPELQKPFFK-OH (KLH) peptide, synthesized by Research Genetics, Huntsville, AL.

Deposit Information:

The following materials were deposited with the American Type Culture Collection:

<u>Name</u>	<u>Deposit Date</u>	<u>Accession No.</u>
<i>Escherichia coli</i> 5 α FGF98	February 26, 1998	98662
<i>Escherichia coli</i> DH5- α FGF8h#4	February 18, 1998	98677

The above materials have been deposited with the American Type Culture Collection, 10801 University Blvd., Manassas, VA 20110-2209, telephone 703-365-2700, under the accession numbers indicated. This deposit will be maintained under the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for purposes of Patent Procedure. The deposits will be maintained for a period of 30 years following issuance of this patent, or for the enforceable life of the patent, whichever is greater. Upon issuance of the patent, the deposits will be available to the public from the ATCC without restriction.

These deposits are provided merely as convenience to those of skill in the art, and are not an admission that a deposit is required under 35 U.S.C. § 112. The sequence of the polynucleotides contained within the deposited materials, as well as the amino acid

• • •

2

2

STATION	DATE	TIME	WIND	TEMP	REL. HUM.	SEA	REMARKS
1	1950	0000	000	50.0	85	0	Cloudy
2	1950	0100	000	50.0	85	0	Cloudy
3	1950	0200	000	50.0	85	0	Cloudy
4	1950	0300	000	50.0	85	0	Cloudy
5	1950	0400	000	50.0	85	0	Cloudy
6	1950	0500	000	50.0	85	0	Cloudy
7	1950	0600	000	50.0	85	0	Cloudy
8	1950	0700	000	50.0	85	0	Cloudy
9	1950	0800	000	50.0	85	0	Cloudy
10	1950	0900	000	50.0	85	0	Cloudy
11	1950	1000	000	50.0	85	0	Cloudy
12	1950	1100	000	50.0	85	0	Cloudy
13	1950	1200	000	50.0	85	0	Cloudy
14	1950	1300	000	50.0	85	0	Cloudy
15	1950	1400	000	50.0	85	0	Cloudy
16	1950	1500	000	50.0	85	0	Cloudy
17	1950	1600	000	50.0	85	0	Cloudy
18	1950	1700	000	50.0	85	0	Cloudy
19	1950	1800	000	50.0	85	0	Cloudy
20	1950	1900	000	50.0	85	0	Cloudy
21	1950	2000	000	50.0	85	0	Cloudy
22	1950	2100	000	50.0	85	0	Cloudy
23	1950	2200	000	50.0	85	0	Cloudy
24	1950	2300	000	50.0	85	0	Cloudy

POLYNUCLEOTIDE AND POLYPEPTIDE SEQUENCES.

SEQ ID NO:1 is the nucleotide sequence.

The reading frame starts at nucleotide 610

CTATAGAAGGTACGCTGCAGGTACCGGTCCGGAATTCCTGGGTCGACCCACGCGTCCGCCACGCGTCCGCGGACGCGTGGGGACGCG
GAGGAGGAGACATGAGCCGGCGGGCGCCAGACGGAGCGGCGTGACGCTTTCGCGCTGCAGCCGCGCGCCCGACCCCGAGCGCTGA
CCCCTGCCCCACGACGCTCCGCGCCGGGCGGAGAGCGCAACTCGGCTTCCAGACCCGCGCGCATGCTGTCCCGGACTGAGCCGG
GCAGCCAGCCTCCACGGACGCCCCGACGGCGGGCGGCGGCGAGTGCAGGAGCTTCCCCGACCGGCCAGGCGCTTCTGCACAACGG
CTGCGCCCCCGACGCCCCGCGCCAGCCGGAGGGCGAGCGCTCGGGAGGAGCGCGCGGGGCGCTGATGCCGACGGGCGCGCGCGG
AGCGCCCCGGAGCAGCAGAGTCTGCAGCAGCAGCAGCGGCGAGGAGGAGCAGCAGCAGCGGCGGCGGCGGCGGCGGCGGCGGAG
GCGCCCCGTCCCGCGCGCGGAGCGGACATGTGACGGCTGGGCTAGGAGCGCGCGCTCCCTCCCGCCAGCGATGTATTAGCGCCC
TCCGCTGCACCTGCCTGTGTTTACACTTCTGCTGCTGTGCTTCCAGGTACAGGTGCTGGTTGCCGAGGAGAAGCTGGACTTCCGCAT
CCAGTGGAGAACCAGACGCGGGCTCGGGACGATGTGAGCCGTAAGCAGCTGCGGCTGTACCAGCTCTACAGCCGGACCAGTGGGAAAC
ACATCCAGGTCTTGGGCGCAGGATCAGTGCCCGCGCGGAGGATGGGACAAGTATGCCAGCTCCTAGTGGAGACAGACCTTCGGT
AGTCAAGTCCGATCAAGGGCAAGGAGACGGAATTCTACCTGTGATGAACCGCAAAGGCAAGCTCGTGGGAAGCCGATGGCACCAG
CAAGGAGTGTGTTCATCGAGAAGTTCTGGAGAACAACACACGGCCCTGATGTCGGCTAAGTACTCCGGCTGGTACGTGGGCTTCA
CCAAGAAGGGGCGGCGCTCTAGAGGATCCCCTGAGGGGCCCAAGCTTACGCGTGCATGC

SEQ ID NO:2 is the amino acid sequence

MYAPSACTCLCLHFLLLCFQVQVLVAENVDFRIHVENQTRADDVSRQLRLYQLYSRTSGKHIQVLGRRISARGEDGDKYAQLLVE
TDTFGSQVRIKGKETEFYLCMNRKGKLVGKPDGTSKECVFIEKVLNNYTALMSAKYSGWYVGFTKKGRPL

SEQ ID NO:3 is the primer sequence

GCGAGGATGGGGACAAGTAT

SEQ ID NO:4 is the FGF coding region with the native human 5' and 3' untranslated region

CCCACGCGTCCGCGGACGCGTGGGGACGCGGAGGAGAGACATGAGCCGGCGGGCGCCAGACGGAGCGG
CCGTGACGCTTTCGCGCTGCAGCCGCGCGCCCCGACCCGGAGCGCTGACCCCTGGCCCCACGACGCTCC
GCGCCCGGGCGGAGAGCGCAACTCGGCTTCCAGACCCGCGCGCATGCTGTCCCGGACTGAGCCGGG
AGCCAGCCTCCACGGACGCCCCGACGGCGGGCGGCGAGTGCAGGAGCTTCCCGCACCGGCCAGG
CGCTCTCTGCACAACGGCTGCCGCCCGCAGCCCTGCGCCAGCCGGAGGGCGCAGCGCTCGGGAGGAG
CCGCGCGGGGCGCTGATGCCGAGGGCGCGCGGAGCGCCCCGGAGCAGCAGAGTCTGCAGCAGCAGC
AGCCGGCGAGGAGGAGCAGCAGCAGCGCGCGGCGGCGGCGGCGGCGGCGGAGGCGCCCGGTCCCGG
CGCGCGGAGCGGACATGTGACGGCTGGGCTAGGAGCGCGCGCTCCCTCCCGCCAGCGATGTATTAGC
GCCCTCCGCTGCACCTTGCTGTGTTTACACTTCTGCTGCTGTGCTTCCAGGTACAGGTGCTGGTTGCC
GAGGAGAACGTGGACTTCCGATCCACGTGGAGAACCAGACGCGGGCTCGGGACGATGTGAGCCGTAAGC

AGCTGCGGCTGTACCAGCTCTACAGCCGGACCAGTGGGAAACATCCAGGTCTTGGGCCGAGGATCAG
 TGCCCGCGGGCAGGATGGGGACAAGTATGCCAGCTCCTAGTGGAGACAGACACCTTCGGTAGTCAAGTC
 CGGATCAAGGGCAAGGAGACGGAAATCTACCTGTGCATGAACCGCAAAGGCAAGCTCGTGGGGAAGCCCG
 ATGGCACCAGCAAGGAGTGTGTTCATCGAGAAGGTTCTGGAGAACAACACACGGCCCTGATGTCGGC
 TAAGTACTCCGGCTGGTACGTGGGCTTCACCAAGAAGGGCGGCCGGAAGGGCCCCAAGACCCGGGAG
 AACCAGCAGGACGTGCATTTTCATGAAGCGCTACCCAAGGGGCAGCCGGAGCTTCAGAAGCCCTTCAAGT
 ACACGACGGTGACCAAGAGGTCCCGTCGGATCCGGCCACACACCTGCCTAGGCCACCCCGCCGCGGCC
 CCTCAGGTCGCCCTGGCCACACTCACACTCCAGAAAACATGCATCAGAGGAATATTTTACATGAAAAAT
 AAGGAAGAAGCTCTATTTTTGTACATTGTGTTTAAAGAAGACAAAACTGAACCAAACTCTTGGGGGG
 AGGGGTGATAAGGATTTTATTGTTGACTTGAAACCCCGATGACAAAAGACTCACGCAAGGGACTGTAG
 TCAACCCACAGGTGCTTGTCTCTCTAGGAACAGACAACCTAAACTCGTCCCAGAGGAGGACTTGAA
 TGAGGAAACCAACACTTTGAGAAGCCAAAGTCTTTTTCCCAAAGGTTCTGAAAGGAAAAAAAAAAC
 AAAAAAAAAAAAAAAAAAAAAAAAAA

SEQ ID NO: 5 is the full length amino acid sequence of FGF

MYSAPSACTCLCLHFLLLCFQVQVLVAEENVDFRIHVENQTRARDDVSRKQLRLYQLYSRTSGKHIQVLG
 RRISARGEDGDKYAQLLVETDTFGSQVRIGKETEFYLCMNRKGKLVGKPDGTSKECVFIEKVLNNYTA
 LMSAKYSGWYVGFTKKGRPRKGPKTRENQDVFHMKRYPKQPELQKPFKYTTVTKRSRRIRPHTPA

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118